


Global seizures of ecstasy-type substances^(a): 2004-2014




Year	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
Kg equivalents ^(b)	12,721	9,729	9,782	16,580	6,259	4,937	4,660	3,640	4,944	4,209	9,353


^(a) Includes substances believed to be ecstasy (e.g. MDMA, MDA, MDE) which may not have been confirmed by forensic testing.

^(b) This quantity reflects the bulk weight of seizures, with no adjustment for purity. Seizures reported by volume (litres, millilitres, etc.) are converted to kg equivalents by assuming a weight of 1 kg per litre. Seizures of amphetamines-group substances reported in tablets or similar units are converted using assumed bulk tablet weights between 90mg and 300 mg, depending on the region and specific drug type, and based on information currently available to UNODC. Further details on conversion factors are given in the methodology section accompanying the World Drug Report.

SEIZURES OF ECSTASY-TYPE SUBSTANCES^(a) - as percentage of world total and in kg equivalents^(b) - HIGHEST RANKING COUNTRIES - 2014



SEIZURES OF ECSTASY-TYPE SUBSTANCES BY REGION - as percentage of world total and in kg equivalents^(a) - 2014




^(a) Includes substances believed to be ecstasy (e.g. MDMA, MDA, MDE) which may not have been confirmed by forensic testing.


^(b) This quantity reflects the bulk weight of seizures, with no adjustment for purity. Seizures reported by volume (litres, millilitres, etc.) are converted to kg equivalents by assuming a weight of 1 kg per litre. Seizures of amphetamines-group substances reported in tablets or similar units are converted using assumed bulk tablet weights between 90mg and 300 mg, depending on the region and specific drug type, and based on information currently available to UNODC. Further details on conversion factors are given in the methodology section accompanying the World Drug Report.

^(c) Financial year 2013/14.


**ECSTASY-TYPE SUBSTANCES*
INTERCEPTED - WORLD: 2004 - 2014**




**ECSTASY-TYPE SUBSTANCES*
INTERCEPTED - ASIA: 2004 - 2014**




ECSTASY-TYPE SUBSTANCES* INTERCEPTED - AMERICAS: 2004 - 2014




**ECSTASY-TYPE SUBSTANCES*
INTERCEPTED - EUROPE: 2004 - 2014**



**ECSTASY-TYPE SUBSTANCES*
INTERCEPTED - AFRICA: 2004 - 2014**



**ECSTASY-TYPE SUBSTANCES*
INTERCEPTED - OCEANIA: 2004 - 2014**



* Includes substances believed to be ecstasy (e.g. MDMA, MDA, MDE) which may not have been confirmed by forensic testing.

** This quantity reflects the bulk weight of seizures, with no adjustment for purity. Seizures reported by volume (litres, millilitres, etc.) are converted to kg equivalents by assuming a weight of 1 kg per litre. Seizures of amphetamines-group substances reported in tablets or similar units are converted using assumed bulk tablet weights between 90mg and 300 mg, depending on the region and specific drug type, and based on information currently available to UNODC. Further details on conversion factors are given in the methodology section accompanying the World Drug Report.